

containing aqueous medium in which the acid is present in the internal and external liposome phases; or

- (2) an aqueous medium containing a base which is substantially impermeable through the vesicle to give an basic liposome-containing aqueous medium in which the base is present in the internal and external liposome phases;

(b) adding:

- (1) to the thus-obtained acid liposome-containing aqueous medium a permanently charged, chargeable, or pH titratable chemical species which is a cationic chemical species, or
- (2) to the thus-obtained acid liposome-containing aqueous medium a permanently charged, chargeable, or pH titratable chemical species which is an anionic chemical species; and

(c) adding to the external liposome phase:

- (1) a base to thereby induce the cationic chemical species to pass into the liposomes' internal acidic aqueous phase, or
- (2) an acid to thereby induce the anionic chemical species to pass into the liposomes' internal basic aqueous phase;

wherein said cationic chemical species or said anionic chemical species is accumulated and entrapped within said liposome to produce a stable liposome vesicle-entrapped chemical species, said stability being such that after administration to an animal the chemical species is carried to its destination by the liposome vesicle before significant leakage occurs.

Sub  
G1  
cont

E1  
conced

52. (Amended) A method of preparing a stable liposome vesicle-entrapped chemical species, which method comprises:

(a) forming liposomes in:

- (1) an aqueous medium containing an acid which is substantially impermeable through the vesicle to give an acidic liposome-containing aqueous medium in which the acid is present in the internal and external liposome phases; or
- (2) an aqueous medium containing a base which is substantially impermeable through the vesicle to give a basic liposome-containing aqueous medium in which the base is present in the internal and external liposome phases;

(b) adding:

- (1) to the thus-obtained acid liposome-containing aqueous medium a permanently charged, chargeable, or pH titratable chemical species which is a cationic chemical species, or
- (2) to the thus-obtained acid liposome-containing aqueous medium a permanently charged, chargeable, or pH titratable chemical species which is an anionic chemical species; and

(c) adding to the external liposome phase:

- (1) a base in an amount effective to create a pH gradient between the external liposome phase and the internal liposome phase to

C2  
Sub  
82

*E 2  
conced* *JLB  
A2  
cont* (2)

thereby induce the cationic chemical species to pass into the  
liposomes' internal acidic aqueous phase, or  
an acid in an amount effective to create a pH gradient between  
the external liposome phase and the internal liposome phase to  
thereby induce the anionic chemical species to pass into the  
liposomes' internal basic aqueous phase;

wherein said cationic chemical species or said anionic chemical species is accumulated and  
entrapped within said liposome to produce a stable liposome vesicle-entrapped chemical  
species, said stability being such that after administration to an animal the chemical species  
is carried to its destination by the liposome vesicle before significant leakage occurs.

---